Staying Healthy: Recommendations for WOMEN



The Veterans Health Administration (VHA) aims to help you stay healthy. The table below lists the **preventive health services** (screening tests, medications, health counseling, and vaccines) that VHA recommends. Screening tests are used to look for health conditions before there are symptoms. These recommendations apply only to adult women of **average risk**. You are a woman of average risk if you have **no personal or family history or symptoms** of the conditions listed below. If you are having symptoms of a condition, please talk with your provider.

Which preventive health services are recommended for average risk women in my age group?

Green: Recommended	Yellow: Recommer	nded for some wome	en – talk with your _l	provider Rec	l: Not recommended	(NR)		
SCREENING TESTS FO	OR AVERAGE	RISK VETERAI	NS					
HEALTH CONDITIONS	18–29 years	30–39 years	40–49 years	50–59 years	60–69 years	70–79 ye	years 80 years and older	
Breast Cancer	Not recommended	Talk with your						
Service Act Veterans	(age 18–29)	provider (age 30-39) Requests for annual screening can be honored after discussion of risks and benefits (age 75 and older)						
Cervical Cancer	Every 3 years (age 21–29)	Recommended. Every 3 or 5 years, depending on tests chosen (age 30–65) Not recommended for those with prior adequat (age 66 and older)						screening
Colon Cancer	Not reco	mmended (age 18–44)		Recommended. Frequency varies by test chosen (age 45–75)			Talk with your provider (age 76–85) NR (age	
Depression				Recommended every ye	ar			
Hepatitis B Infection			Recommended for sor	ne women – talk with your	provider (age 18 and older)			
Hepatitis C Infection		Reco	mmended in adults aged	18-79 years, and in all preg	nant women, during each pr	regnancy		
High Blood Pressure	Recommended every year (age 18 and older)							
HIV Infection		Recommende	ed once (age 18 and older); in all pregnant Veterans;	annually in Veterans with on	going risk factor	rs	
Lung Cancer	Not recommended (age 18–49) Recommended for some women – talk with your provider (age 50–80) NR (age 81 and older							
Osteoporosis	Recommended for some women — talk with your provider (age 18—64) Recommended for some women — talk with your provider (age 18—64) Recommended for some women — talk with your provider (age 18—64) Recommended for some women — talk with your provider (age 18—64)							
Sexually Transmitted Infections	Recommended. Test for gonorrhea and chlamydia (age 18–24). Talk with your provider about syphilis (age 25 and older) Recommended for some women, talk with your provider about testing for gonorrhea, chlamydia, and syphilis (age 25 and older) Recommended for some women, talk with your provider about testing for gonorrhea, chlamydia, and syphilis (age 25 and older) Recommended for all pregnant Veterans, early screening for syphilis infection, regardless of age.							
MEDICATIONS								
HEALTH CONDITIONS	18–29 years	30–39 years	40–49 years	50–59 years	60–69 years	70–79 ye	ars 80 years	s and olde
Folic Acid for Pregnancy Planning		ed. Daily folic acid suppleme become pregnant (ac	ent for any woman who n				age (age 55 and older)	, and order
Aspirin to Prevent Cardiovascular Disease	Not recommen	ded (age 18–39)	rovider (age 40–59)	der (age 40–59) Not recommended (age 60 and older)				
PrEP to Prevent HIV Infection	Talk with your provider (age 19 and older)							
Statin to Prevent Cardiovascular Disease	N	ot recommended (age 19–4		Recommended for some women — talk with your provider (age 40 and older)				
HEALTH COUNSELIN	 G			_				
HEALTH CONDITIONS	18–29 years	30–39 years	40–49 years	50–59 years	60–69 years	70–79 ye	ars 80 years	s and olde
Tobacco Use	Recommended every visit (if using tobacco) (age 18 and older)							
Alcohol Use	Talk with your provider about healthy alcohol use (age 18 and older)							
Healthy Diet and Physical Activity	Talk with your provider about a healthy diet and physical activity (age 18 and older)							
Overweight & Obesity	Recommended every year (age 18 and older)							
/ACCINES						T		
HEALTH CONDITIONS	19–29 years	30–39 years	40–49 years	50–59 years	60–69 years	70–79 ye		s and olde
COVID-19	Re	ecommended for all women			o-date (<u>https://www.va.gov</u>	<u>/health-care/co</u>	<u>vid-19-vaccine/</u>)	
Hepatitis A				<u> </u>	provider (age 19 and older)			
Hepatitis B	Recommended for all women who have not already had Hepatitis B vaccine (age 19-59) Recommended for some women - talk with your provider (age 60 and old							
Herpes Zoster (Shingles) RZV vaccine	Recommended for some women — talk with your provider (age 19 and older)			Recommended	Recommended 2 doses (age 50 and older), with second dose 2—6 months after first dose			
Human Papilloma Virus (HPV) HPV9 vaccine	2–3 doses (age 19–26) if series not completed	Talk with your provider (ag	je 27–45)		Not recommended (age	46 and older)		
Influenza (Flu) Injectable, Inhaled			Recon	nmended every year (age 19	and older)			
Measles, Mumps, and Rubella (MMR)	Recommended for some women — talk with your provider (age 19 and older)							
Meningococcal Disease	Recommended for some women — talk with your provider (age 19 and older)							
Pneumococcal Disease PCV21 vaccine, PCV20 vaccine, PCV15 vaccine, PPSV23 vaccine	Recommended for some women — talk with your provider (age 19—49) Recommended - talk with your provider to make sure you are up-to-date (age 50 and older)							
Respiratory Syncytial Virus (RSV) RSV vaccine	Pregnant Veterans are recommended to receive a one-time (single lifetime) RSV Abrysvo® vaccine between 32-36 weeks gestation during RSV season "One time" (single lifetime) dose recommended for some women – talk with your provider (age 60–74) "One time" (single lifetime) dose recommended for some women – talk with your provider (age 60–74)							
Tdap (Tetanus, Diphtheria		(Or for wou		st 1 dose. Get additional do	se with every pregnancy	ostor vaccino)		



& Pertussis)

Td (Tetanus and diphtheria)

Varicella (Chickenpox)





Recommended for some women - talk with your

provider (2 doses if born 1980 or later)

(Or for wound management if > 5 years have passed since last tetanus toxoid-containing booster vaccine).

Recommended. Booster every 10 years (age 19 and older) (Either Tdap or Td may be used)

(Or for wound management if > 5 years have passed since last tetanus toxoid-containing booster vaccine).